

*SwissMountain* has a major advantage compared to most bottled waters ie

It contains virtually no harmful Nitrate (below 0.1 mg/l)

To help consumers understand the importance of low levels of nitrate we provide below excerpts from a Cornell University paper.

### **Nitrate: Health Effects in Drinking Water**

by

Margaret McCasland, Nancy M. Trautmann, and Keith S. Porter  
Center for Environmental Research  
and  
Robert J. Wagenet  
Dept. of Agronomy  
Cornell University

**Nitrate is one of the most common groundwater contaminants** in rural areas. It is regulated in drinking water primarily because excess levels can cause methemoglobinemia, or "blue baby" disease. Although nitrate levels that affect infants do not pose a direct threat to older children and adults, they do indicate the possible presence of other more serious residential or agricultural contaminants, such as bacteria or pesticides.

**Nitrate in groundwater originates primarily from fertilizers, septic systems, and manure storage or spreading operations.**

#### **What Is Nitrate?**

Nitrate is an inorganic compound that occurs under a variety of conditions in the environment, both naturally and synthetically. Nitrate is composed of one atom of nitrogen (N) and three atoms of oxygen (O); the chemical symbol for nitrate is NO<sub>3</sub>. Nitrite (NO<sub>2</sub>) can be formed from nitrate by a chemical process called reduction. Nitrate does not normally cause health problems unless it is reduced to nitrite.

Nitrate in drinking water is measured either in terms of the amount of nitrogen present or in terms of both nitrogen and oxygen. The federal standard for nitrate in drinking water is 10 milligrams per liter (10 mg/l) nitrate-N, or 45 mg/l nitrate-NO<sub>3</sub>. when the oxygen is measured as well as the

nitrogen. Unless otherwise specified, nitrate levels usually refer only to the amount of nitrogen present, and the usual standard, therefore, is 10 mg/l.

**Short-term exposure to drinking water with a nitrate level at or just above the health standard of 10 mg/l nitrate-N is a potential health problem primarily for infants.** Babies consume large quantities of water relative to their body weight, especially if water is used to mix powdered or concentrated formulas or juices. Also, their immature digestive systems are more likely than adult digestive tracts to allow the reduction of nitrate to nitrite. In particular, the presence of nitrite in the digestive tract of newborns can lead to a disease called methemoglobinemia.

### **Infant Feeding Practices to Minimize Intake of Nitrate and Nitrite**

1. Breast feeding. Little if any nitrate gets into breast milk, unless the mother is consuming very large quantities of nitrate. Also, bacterial contamination is not a problem when breast milk is consumed directly.
2. Bottle feeding. Use already diluted liquid formulas or use low-nitrate water to dilute concentrated liquid or powdered formulas. Also, mixed formulas should be kept under refrigeration and used promptly to minimize bacterial reduction of nitrate to nitrite.
3. Vegetables. Since many vegetables are high in nitrate, their consumption should be limited until an infant is 4-6 months old and their digestive tract has sufficiently matured. Your physician can help you decide when to add new foods. Vegetables should always be prepared while fresh and refrigerated promptly after cooking to minimize bacterial activity.

### **What Is Methemoglobinemia?**

**Methemoglobinemia is the most significant health problem associated with nitrate in drinking water.** Blood contains an iron-based compound called hemoglobin, which carries oxygen. When nitrite is present, hemoglobin can be converted to methemoglobin, which cannot carry oxygen. In the blood of adults, enzymes continually convert methemoglobin back to hemoglobin, and methemoglobin levels normally do not exceed 1 percent. Newborn infants have lower levels of these enzymes, and their methemoglobin level is usually 1 to 2 percent. Anything above that level is considered methemoglobinemia.

Once diagnosed, methemoglobinemia can be readily reversed, although with anoxia permanent damage may have occurred. Methemoglobinemia can be prevented by restricting consumption of nitrite and nitrate and by limiting the opportunities bacteria have to reduce nitrate in food to nitrite before consumption.

Consuming drinking water with nitrate levels near the drinking water standard does not normally increase the methemoglobin level of humans beyond infancy. Some individuals, however, may have increased susceptibility to methemoglobinemia due to exposure to antioxidant medications and chemicals, or other conditions that may inhibit the body's ability to reconvert methemoglobin to hemoglobin (such as pregnancy or certain rare diseases).

Nitrate in drinking water starts affecting the health of the general populace at levels in the range of 100 to 200 mg/l nitrate-N, but the effect on any given person depends on many factors, including other sources of nitrate and nitrite in the diet. Some of the nitrate consumed can be converted in the body to nitrite, which under appropriate circumstances can combine with amines (portions of protein molecules often found in foods, medications, cigarette smoke, decaying

plants, soil, and sometimes water) to form nitrosamines, well- documented cancer-causing substances. So far, the only studies linking nitrate in drinking water with cancer have involved nitrate levels that are quite high (at or above 100-200 mg/l nitrate-N).

## **How are Safety Standards for Drinking Water Set?**

Congress passed the federal Safe Drinking Water Act in 1974. The U S. Environmental Protection Agency was given responsibility for setting drinking water standards for all the states, and each state became responsible for enforcing these standards. In New York State, the Department of Health regulates all public water supplies serving 25 or more people.

Because potential health risks are often unknown or hard to predict, many drinking water standards are set at some fraction of the level of "no-observed adverse-health effects." In general, the greater the uncertainty about potential health effects, the greater the margin of safety built into the standard.

In the case of nitrate, there may not be a large safety factor. A 1977 report by the National Academy of Science concluded that "available evidence on the occurrence of methemoglobinemia in infants tends to confirm a value near 10 mg/l nitrate as nitrogen as a maximum no-observed adverse-health-effect level, but there is little margin of safety in this value."

## **How Can I Lower the Nitrate Level in My Current Water Supply?**

**There are no simple ways to remove nitrate from water in the home. Because nitrate does not evaporate the way chlorine does, boiling, freezing, or letting water stand does not reduce the nitrate level. In fact, boiling water for more than 10 minutes can make the nitrate more concentrated. Boiling water in an aluminum pan may also convert nitrate to nitrite.**

## **Where Can I Get Low-Nitrate Water?**

**If your private or public water supply has elevated nitrate levels, then purchasing bottled water is one of your best alternatives. Even in bottled water, nitrate levels can vary considerably. Some states require bottled water companies to have their water tested for nitrate annually. As long as the results are below the state standard, the companies don't have to report the actual levels to the Department of Health, but they should be willing to give exact test results to consumers. In general, distilled or mineral-reduced water is more likely to have low-nitrate levels than spring or mineral water.**

Municipal water supplies are another potential source of low nitrate water. City water supplies often come from large lakes or reservoirs where any source of nitrate is likely to be diluted by the large quantity of water.

Hauling your own water from springs, lakes, or streams can be risky, both because of the unknown levels of nitrate and because of the possible presence of other contaminants, especially bacteria. Private wells also present the risk of unknown nitrate levels, but are more likely to have been tested for bacterial contamination.

Risks also are associated with collecting rainwater. Rooftop cisterns often contain lead solder, which can be dissolved by rainwater. Also, rain collected off a roof can contain asbestos or other contaminants from the roofing material.

Alternative water supplies are only as good as their source, their collection method, and their storage. Before using any source of water, find out exactly what the levels of nitrate and bacteria are in that supply. Bacterial contamination is a double problem: it may be responsible for the reduction of nitrate to nitrite; but more importantly, the bacteria themselves can cause illness.

---

**Disclaimer: Please read the pesticide label prior to use. The information contained at this web site is not a substitute for a pesticide label. Trade names used herein are for convenience only; no endorsement of products is intended, nor is criticism of unnamed products implied. Most of this information is historical in nature and may no longer be applicable.**

For more information relative to pesticides and their use, please contact the PMEPP staff at:

5123 Comstock Hall  
Cornell University  
Ithaca, NY 14853-0901  
(607)255-1866



Cornell University